

Supplementary information

The IL-17A rs2275913 single nucleotide polymorphism is associated with protection to tuberculosis but related to higher disease severity in Argentina

Rolandelli, A.^{1,2}; Hernández Del Pino, R. E.^{1,3}; Pellegrini, J. M.^{1,2}; Tateosian, N. L.^{1,2}; Amiano, N.O.^{1,2}; de la Barrera S⁴; Casco, N.⁵; Gutiérrez, M.⁶; Palmero, D. J.⁵; García, V. E.*^{1,2}

¹ Departamento de Química Biológica. Facultad de Ciencias Exactas y Naturales. UBA, Intendente Güiraldes 2160, Pabellón II, 4º piso, Ciudad Universitaria (C1428EGA), Buenos Aires, Argentina.

² Instituto de Química Biológica, Facultad de Ciencias Exactas y Naturales (IQUIBICEN), UBA (Universidad de Buenos Aires)-CONICET, Intendente Güiraldes 2160, Pabellón II, 4º piso, Ciudad Universitaria (C1428EGA), Buenos Aires, Argentina.

³ Centro de Investigaciones y Transferencia del Noroeste de Buenos Aires (CIT NOBA), CONICET. Newbery 261, Junín (6000), Buenos Aires, Argentina.

⁴ Instituto de Medicina Experimental-CONICET-Academia Nacional de Medicina. Pacheco de Melo 3081 (CP1425), Buenos Aires, Argentina.

⁵ División Tisiopneumología Hospital F.J. Muñoz, Uspallata 2272, (C1282AEN) Buenos Aires, Argentina.

⁶ Sección Bacteriología de la Tuberculosis, Hospital General de Agudos “Dr. E. Tornu”, Combatientes de Malvinas 3002, (C1427ARN) Buenos Aires, Argentina


*vgarcia@qb.fcen.uba.ar

Supplementary Table S1.

		HD (N=207)			TB (N=185)		
rs2275913 genotypes		GG	GA	AA	GG	GA	AA
Sex	Male	40	31	10	98	41	6
	Female	68	45	13	26	10	4
<i>P value</i>		0.79			0.34		


Genotypic frequencies of the IL-17A rs2275913 SNP in HD and TB populations stratified by sex. *P* values were calculated by the Chi-Square test for categorical variables. HD: healthy donors; TB: tuberculosis patients.

Supplementary Figure S1



Supplementary Figure S1. IL-17A rs2275913 SNP genotyping by ARMS-PCR method. (a) Primer sequences (two Forward primers and one common Reverse primer) designed to specifically amplify a 317 pb amplicon that discriminates both alleles of the rs2275913 SNP. **(b)** Image of an agarose gel displaying the PCR products obtained from three genotypically different individuals for the SNP under study is shown. PCR positive control: Human Growth Hormone (HGH) gene fragment (440 bp). rs2275913 genotypes were assessed from the presence/absence of PCR amplicon corresponding to the specific allele (A or G). **(c)** DNA sequencing of the amplicons obtained from three genotypically different individuals. Primers specificity of the rs2275913 SNP were confirmed by direct sequencing of the amplified IL-17A gene fragment by Sanger method and a 100% concordance was obtained among the results obtained from ARMS-PCR and DNA sequencing.

Supplementary Figure S2



Supplementary Figure S2. Percentage of IFN- γ ⁺IL-17A⁺CD4⁺T cells in *Mtb*-Ag stimulated PBMCs from HD and TB carrying the genotypic variants of the rs2275913 SNP. PBMCs from HD (n=15, white bars, left panel) and TB (n=16, black bars, right panel) carrying the different genotypes of the rs2275913 SNP were stimulated for five days with *Mtb*-Ag, and IFN- γ ⁺IL-17A⁺CD4⁺T cells percentage was determined by Flow Cytometry. The percentages represent an increase in the number of cytokine-positive CD4⁺T cells in response to *Mtb*-Ag stimulation. IL-17A and IFN- γ expression was determined gating on lymphocytes by light scatter first, and then gating on CD4⁺T cells. Bars represent the Mean \pm SEM. P values were calculated by the Kruskal-Wallis (ANOVA) test for unpaired and non-parametric samples. *P<0.05.